



Dental Implant & Periodontal Surgeons, P.C.

Periodontics, Dental Implants, PiezoSurgery & IV Sedation

Sam Khoury, DDS MS

Diplomate-American Board of Periodontology

Juan C. Cabrera, DMD MS

Diplomate-International Congress of Oral Implantology


Ridge/Jaw Bone Augmentation

A **Ridge augmentation/modification** is a common dental procedure often performed to help recreate the natural contours of the gums and jaw that may have been lost due to bone loss as a result of a tooth extraction, or trauma. The ridge of the jaw is the bone that surrounds the roots of the teeth. When a tooth is removed, an empty socket is left in the alveolar ridge bone. Most of the time, bone graft is added after extraction to preserve the width and height of the jaw. This is called "Socket Preservation." If this procedure is not

performed, the bone surrounding the socket breaks and resorbs. The previous height and width of the socket will continue to deteriorate causing resorption. **Figure 1**

Rebuilding the original height and width of the alveolar ridge is required for dental implant placement, or for aesthetic purposes. Dental implants require bone to support their structure, and a **ridge augmentation** can help rebuild this bone to accommodate the implant.

Figure 1
Resorbed ridge



See next page.

Sam B. Khoury DDS MS™



Practice Limited to Dental Implants & Periodontics & IV Sedation

638 Newtown Yardley Road # 1C Newtown PA 18940 • 215 968 9601 • info@treatmygums.com

107 Commons CT Chadds Ford, PA 19317 • 610 500 0610 • treatmygums@gmail.com

How is the Oral Surgery Accomplished?

During the ridge modification procedure, the gum is lifted away from the ridge to fully expose the defect in the bone. The bony defect can be filled with bone graft material which can help regenerate lost bone. Two types of barriers can be used to contain the bone graft. One can be a resorbable barrier "Figure 2" and the other is a titanium barrier "Figure 3." Finally, the incision is closed and several months of healing will be required. Depending on the case and type of implant and procedure, a dental implant may be placed during the ridge modification procedure or when healing is complete. Ridge modification improves the cosmetic appearance, functionality of the mouth, and the ability to enjoy implants for many years.



Resorbable Membrane

VS

Titanium Membrane

Sam B. Khoury DDS MS™